

WO 00/09557 A1

Job No.: 1505-80958

Translated from Japanese by the Ralph McElroy Translation Company
910 West Avenue, Austin, Texas 78701 USA

INTERNATIONAL PATENT OFFICE
WORLD ORGANIZATION FOR INTELLECTUAL PROPERTY

International patent published on
the basis of the Patent Cooperation Treaty
INTERNATIONAL PUBLICATION NO. WO 00/09557 A1

International Patent Classification ⁶ :	C 07 K	14/47
	C 12 N	15/12
		//5/10
	C 12 P	21/02
	(C 12 P	21/02
	C 12 R	1:91)
International Filing No.:	PCT/JP99/04352	
International Filing Date:	August 11, 1999	
International Publication Date:	February 24, 2000	
Priority		
Date:	August 12, 1998	
Country:	JP	
No.:	Hei 10[1998]-227723	
Designated States:	AU, CA, CN, KR, US, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE)	

NOVEL GENE AND PGTH PROTEIN ENCODED THEREBY

Inventors and	
Inventors/Applicants (only for US):	Osamu Ohara; Takahiro Nagase; Nobuo Nomura Kazusa DNA Research Institute Foundation 1532-3 Yakuni, Kisarazu-shi, Chiba-ken (JP)

Kiyoshi Takayama
Hitoshi Toyoda;
Makoto Yoshimoto
Taisho Pharmaceutical Co., Ltd.
3-24-1 Takata, Toshima-ku,
Tokyo-to (JP)

Applicant (for all designated
states other than US):

Kazusa DNA Research Institute
Foundation
1532-3 Yakuni
Kisarazu-shi, Chiba-ken (JP)

Taisho Pharmaceutical Co., Ltd.
3-24-1 Takata, Toshima-ku,
Tokyo-to (JP)

Agent:

Tomizo Kitagawa
Taisho Seiyaku K.K.
3-24-1 Takata, Toshima-ku,
Tokyo-to (JP)

Published
With International Search Report.

FOR INFORMATION ONLY

Codes for the identification of PCT contract states on the cover sheets of the documents that publish the international applications in accordance with the PCT.

AE	United Arab Emirates	KP	Democratic People's	YU	Yugoslavia
AL	Albania		Republic of Korea	ZA	South Africa
AM	Armenia	KR	South Korea	ZW	Zimbabwe
AT	Austria	KZ	Kazakhstan		
AU	Australia	LC	Saint Lucia		
AZ	Azerbaijan	LI	Liechtenstein		
BA	Bosnia-Herzegovina	LK	Sri Lanka		
BB	Barbados	LR	Liberia		
BE	Belgium	LS	Lesotho		
BF	Burkina Faso	LT	Lithuania		
BG	Bulgaria	LU	Luxembourg		
BJ	Benin	LV	Latvia		
BR	Brazil	MA	Morocco		
BY	Belarus	MC	Monaco		
CA	Canada	MD	Republic of Moldavia		
CF	Central African Republic	MG	Madagascar		
CG	Congo	MK	Macedonia (former Yugoslavian Republic of Macedonia)		
CH	Switzerland	ML	Mali		
CI	Côte d'Ivoire	MN	Mongolia		
CM	Cameroon	MR	Mauritania		
CN	China	MW	Malawi		
CR	Costa Rica	MX	Mexico		
CU	Cuba	NE	Niger		
CY	Cyprus	NL	Netherlands		
CZ	Czech Republic	NO	Norway		
DE	Germany	NZ	New Zealand		
DK	Denmark	PL	Poland		
DM	Dominica	PT	Portugal		
EE	Estonia	RO	Romania		
ES	Spain	RU	Russian Federation		
FI	Finland	SD	Sudan		
FR	France	SE	Sweden		
GA	Gabon	SG	Singapore		
GB	United Kingdom	SI	Slovenia		
GD	Grenada	SK	Slovakia		
GE	Georgia	SL	Sierra Leone		
GH	Ghana	SN	Senegal		
GM	Gambia	SZ	Swaziland		
GN	Guinea	TD	Chad		
GR	Greece	TG	Togo		
HR	Croatia	TJ	Tajikistan		
HU	Hungary	TM	Turkmenistan		
ID	Indonesia	TR	Turkey		
IE	Ireland	TT	Trinidad and Tobago		
IL	Israel	TZ	Tanzania		
IN	India	UA	Ukraine		
IS	Iceland	UG	Uganda		
IT	Italy	US	United States of America		
JP	Japan	UZ	Uzbekistan		
KE	Kenya	VN	Vietnam		
KG	Kyrgyzstan				

Field of the technology

This invention pertains to a novel PGTH protein of human brain origin having a prostaglandin transport activity and the pgth gene encoding the protein.

Prior art

Prostaglandin is a generic name for a series of physiologically active lipids such as prostaglandin E, prostaglandin D, prostaglandin F, prostaglandin I, prostaglandin J, etc. Prostaglandin is a physiologically active substance inside the body strongly related to control of physiological functions such as blood flow rate, sleeping, gastric mucosa protective action, thrombus formation, pregnancy, etc., through specific cell membrane or intranuclear receptors.

Prostaglandin is produced inside cells as a result of eicosapolyenic acids such as arachidonic acid, etc., being cut out by phospholipase A2 from the cell membrane and converted with cyclooxygenase and various prostaglandin synthetic enzymes by responding to various physiological stimuli, and after being released outside the cells, it has autocrine or paracrine effects. On the other hand, liberated prostaglandin is also circulated in the blood flow, taken up by a specific cell, metabolized and so disappears.

A trace amount of prostaglandin shows a strong physiological activity, and consequently, the production of prostaglandin compounds is strictly controlled by controlling the activity of production-related and metabolism-related enzymes.

However, prostaglandin has been reported to be unable to pass through the lipid double layer of the cell membrane by itself. Therefore, as a prostaglandin transport mechanism, the presence of a special protein has been presumed in the process of prostaglandin produced inside a cell exiting it and the process of prostaglandin circulating in the blood flow being taken up into a specific cell.

As a protein involved in the transport mechanism described above, prostaglandin transporter (abbreviated hPGT: human prostaglandin transporter, below) has been reported, but it is not a protein involved in the transport of all prostaglandin compounds, and there are many unclear points. Consequently, it is thought that if a biological molecule other than hPGT involved in the transport mechanism can be elucidated, the biological molecule found might be usable directly as a medical drug or indirectly as a compound for studying compounds that might be usable as a medical drug. Therefore, the objective of this invention is to identify such a molecule and use it as a medical drug or for the development of medical drugs.

Presentation of the invention

The inventors of this invention studied diligently to find the desired protein by using genes expressed in the human brain, and as a result, they found the presence of a novel PGTH protein (prostaglandin transporter homologue), successfully isolated a pgth gene encoding the protein, and they arrived at this invention.

Specifically, this invention pertains to (a) a protein having the amino acid sequence described in sequence No. 1 or (b) a protein having an amino acid sequence with 1 to several amino acids deleted, substituted or added to the amino acid sequence of sequence No. 1, and having a prostaglandin transport activity.

Furthermore, this invention also pertains to (c) a gene comprising DNA described sequence No. 2 or (d) DNA which can be hybridized with the DNA of sequence No. 2 under stringent conditions and which encodes a protein having a prostaglandin transport activity.

The pgth gene of this invention can be isolated as a cDNA fragment containing the gene from a cDNA library of human brain origin. The cDNA library used by the inventors of this invention was prepared based on commercially available mRNA of human brain origin from the Clontech Co.

As a method for identifying the cDNA encoding a protein having a prostaglandin transport activity in the cDNA library described above, the method of Ohara, et al., (DNA Research 4: p 53, 1997) was used as an extensive cDNA library analysis method using a long-chain cDNA library. From a long-chain cDNA library of human brain origin prepared by the method of Ohara, et al., 25,000 recombinants are randomly selected, the 5' and 3' - base sequences of the cDNA from 15,000 clones were determined, and a clone showing homology to the gene encoding hPGT already reported from the 5' sequences of all the clones can be found by using a DNA analysis program (BLAST and FastA).

The presence of a region encoding the protein (ORF: open reading frame) in the base sequence can be confirmed by a conventional method using a computer program. After becoming confident of the presence of the desired gene in the cDNA sequence, the inventors of this invention found one ORF in the sequence by utilizing a computer, the gene was named pgth, and the protein encoded by the gene was named PGTH. The PGTH of the invention is a protein comprising a total of 709 amino acid residues and having a molecular weight of about 80 kd.

The invention pgth is a gene comprising 2130 bp shown in sequence No. 2. By using this pgth and conventional genetic recombination techniques using a suitable host vector system, it is possible to prepare a recombinant gene. As a suitable vector, there are plasmids of *E. coli* origin (such as pBR322, pUC118, etc.), of *Bacillus subtilis* origin, (such as pSH19, etc.) yeast origin plasmid (such as pUB110, pC194, etc.), bacteriophages, animal viruses such as retroviruses,

vaccinia virus, etc., etc. At the time of recombination, it is possible to add translation initiation and termination codons using suitable DNA adaptors. Furthermore, for gene expression, a suitable expression promoter is attached upstream of the gene. The promoter to be used is suitably selected depending on the host used. For example, if the host is *E. coli*, there are T7, lac, trp, λ PL promoters, etc.; if the host is a *Bacillus*, there are SPO promoters, etc.; if the host is a yeast, there are PHO5, GAP, ADH promoters, etc.; and if the host is an animal cell, there are SV40-origin, retrovirus promoters, etc.

Furthermore, the gene may be expressed as a fused protein with another protein (such as glutathione-S-transferase, protein A, etc.). In the case of a fused PGTH prepared by using such a method, a suitable protease (such as thrombin, etc.), may be used to cut out the protein.

As a host usable in the case of PGTH expression, there are various strains of *Escherichia coli*, various strains of *Bacillus subtilis*, various strains of the yeast *Saccharomyces cerevisiae* and animal cells such as COS-7, CHO cells, etc.

As a method for transforming a host cell using the above recombinant vector, a specific method conventionally used to transform the selected host cell is used.

Incidentally, in this invention, DNA which has a DNA sequence other than that shown in sequence No. 2 which can be hybridized with the DNA and encodes a protein having a prostaglandin transport activity, is also included in the scope of this invention.

Specifically, DNA which has a DNA sequence, the total length of the pgth sequence, partially changed due to various artificial treatments such as random mutations, introduction of site-specific mutations, or mutagen treatment, DNA fragment mutation, deletion ligation after scission with restriction enzymes, is also included in the scope of this invention in spite of having a DNA sequence different from that of sequence No. 2 as long as such a DNA variant can be hybridized with pgth under stringent conditions and encodes a protein having a prostaglandin transport activity.

The extent of the above DNA mutation is within the allowable range if the variant has 90% or higher homology with the DNA sequence of pgth. Furthermore, as an extent of hybridization with pgth, Southern hybridization with pgth may be carried out under conventional conditions, for example, in the case of probe labeling with a DIG DNA Labeling kit (Boehringer-Mannheim Cat. No. 1175033), hybridization conditions of a DIG Easy Hyb solution (Boehringer-Mannheim Cat. No. 1603558) at 32°C and washing of the membrane in a 5X SSC solution (containing 0.1% w/v SDS) at 50°C (1X SSC comprises 0.15M NaCl and 0.015M sodium citrate).

Furthermore, a protein encoded by the gene variant which is highly homologous to pgth as described above and has a prostaglandin transport activity is also included in the scope of this invention.

Specifically, a variant having one or more amino acids deleted, substituted or added to the amino acid sequence of PGTH is included in the scope of this invention as long as this variant is a protein having a prostaglandin transport activity.

The side chains of the amino acids, which are the constituent elements of proteins are respectively different with respect to hydrophobicity, electrical charge, size, etc., but several highly conservative relationships in the meaning of practically not affecting the three-dimensional structure (it is also called the steric structure) of proteins have been known from experiences or actual physicochemical observations. For example, for substitution of amino acid residues, there are glycine (Gly) and proline (Pro), Gly and alanine (Ala) or valine (Val), leucine (Leu) and isoleucine (Ile), glutamic acid (Glu) and glutamine (Gln), aspartic acid (Asp) and asparagine (Asn), cysteine (Cys) and threonine (Thr), Thr and serine (Ser) or Ala, lysine (Lys) and arginine (Arg), etc.

Therefore, any variant protein due to substitution, insertion, deletion, etc., in the amino acid sequence of the PGTH shown in sequence No. 1 can be said to be within the scope of this invention if the variation is a variation which conserves the three-dimensional structure of the PGTH, and the protein is a protein having a prostaglandin transport activity similar to PGTH. The allowable extent of this variation is 90% or higher homology with the amino acid sequence shown in sequence No. 1.

Industrial application field

The abnormal expression of pgth or functional failure of PGTH is presumed to be a critical disorder because PGTH has a prostaglandin transport activity, and consequently the normal prostaglandin production mechanism of the body is lost.

Therefore, PGTH itself is considered to be useful as a drug, and on the other hand, pgth or PGTH may be used for effectively studying or evaluating a substance having the same function as that of PGTH, a substance promoting or inhibiting its function, a substance promoting the expression of the gene, etc.

Best embodiment of the present invention

This invention is explained further in detail using application examples as follows, but this invention is certainly not limited to these application examples. Incidentally, unless specified, the experimental procedures used in the following application examples are those

described in standard experimental manuals such as *Molecular Cloning*, 2nd ed. (Cold Spring Harbor Laboratory Press, 1989), etc., and the operating manuals in commercially available kits, and they can be carried out under the conditions recommended for the respective commercially available products such as restriction enzymes, etc.

Application Example 1 Cloning of pgth

1) Construction of a long chain cDNA library of human brain origin

An oligonucleotide (GACTAGTTCTAGATCGCGAGCGGCCGCCC(T)₁₅) containing a NotI site was synthesized using a DNA synthesizer (ABI380B). It was used as a primer, and a double chain cDNA was synthesized using mRNA of human brain origin as a template and the SuperScript II reverse transcriptase kit (Gibco BRL). The ligation of the synthetic DNA was carried out with the cDNA and SalI site-containing adapter (Takara Shuzo), subsequently, NotI digestion was carried out, and cDNA fragments of 3 kb or larger were purified using electrophoresis with a 1% concentration of low-melting agarose.

After ligation of the purified cDNA fragments with a SalI-NotI restriction enzyme-treated pBluescriptII SK⁺ plasmid, the recombinant plasmids were introduced into *E. coli* ElectroMax DH10B strain (Gibco BRL) using the electroporation method. Subsequently, 25,000 recombinants were randomly selected from the library, the recombinant DNAs were extracted, and the 5'- and 3'-base sequences of the cDNAs of 15,000 clones were determined. For the sequence determination, a PE Applied Biosystem Co., DNA sequencer (ABI PRISM377) and the reaction kit from the same company were used.

2) Selection of clones containing the pgth sequence

The 5 sequences of all the clones determined in 1) were compared with the sequence of hPGT already reported using DNA analytical programs (BLAST and FastA), and as a result, a clone named HK07457 showed significant homology.

3) DNA fragment base sequence determination

The base sequence determination was carried out using a PE Applied Biosystem Co. DNA sequencer and the dye primer method. The sequence was mostly determined using the shotgun method, and for a portion of the base sequence, an oligonucleotide was synthesized based on the base sequence already determined, and the primer walking method was used to determine the entire base sequences of the two chains. The entire base sequence of the cDNA of the clone is shown in sequence No. 3.

The cDNA contains an ORF encoding a protein (PGTH) comprising 709 residues. A termination codon was found to appear in the upstream region of a methionine residue, which was an initiation codon of the protein, with the same reading frame. Therefore, the amino acid sequence shown in sequence No. 3 was confirmed to be the only possibility as an amino acid sequence of the protein encoded by the cDNA fragment.

Figure 1 shows the amino acid homology between already reported hPGT and the PGTH of this invention. The two show high homology, especially, the position of the cysteine residue present at the C-terminal of PGTH is preserved, and the 77th residue glutamine, 561st residue arginine and 614th residue lysine of hPGT, which are amino acids especially important for the transport activity, are also preserved in PGTH.

Application Example 2

Confirmation of protein expression by in vitro translation of pgth

The plasmid containing pgth prepared in Application Example 1 was treated with RNase A, subsequently, RNase A was removed using ADVAMAX beads (AGTC Co.), and in vitro translation was carried out using a TNT T7 coupled reticulocyte lysate system (Promega Co.) in the presence of (³⁵S)-methionine. A portion of the reaction mixture was separated by SDS-polyacrylamide gel electrophoresis (SDS-PAGE), and the analysis carried out using BAS-2000 (Fuji Shashin Kogyo). As a result, the presence of a single band at about 80 kd was confirmed as shown in Figure 2.

Application Example 3

Construction of animal cell expression vector

1) Amplification of ORF-containing cDNA

An oligonucleotide (following sequence 1) having a sequence upstream from the initiation codon of the protein of sequence No. 3 and oligonucleotide (following sequence 2) having a sequence of a portion downstream from the termination codon of the protein and the reverse complementary strand chain were synthesized using a DNA synthesizer (ABI Co., Model 380B).

Sequence 1

5-CTGGAGCTCACTGCACTCCAGCAGTC-3

Sequence 2

5-AGCTCACACTCGGGAATCCTCTGGCTTC-3

The recombinant cDNA containing sequence No. 3 isolated in application example 1 was used as a template, the oligonucleotides of the sequences 1 and sequences 2 were used as a primer, and the following PCR procedures were carried out using a Takara LA PCR kit Ver. 2 and the PCR thermal cycler MP (Takara Shuzo).

cDNA	5 μ L (10 ng)
10X PCR buffer (containing 25 mM Mg ⁺⁺)	5 μ L
2.5 mM dNTP	8 μ L
10 μ M Sequence 1	2 μ L
10 μ M Sequence 2	2 μ L
Water	27.5 μ L
LA Taq polymerase	0.5 μ L
Total amount	50 μ L

The PCR cycle was carried out by holding at 94°C for 2 mn, carrying out the reaction at 98°C for 20 sec, cooling to 68°C at a rate of 1°C/2 sec, holding at 68°C for 3 min, at 72°C for 10 min, and repeating 30 times.

The above method was used to amplify a DNA fragment (about 2.2 kb) having a portion of sequence No. 3.

2) Subcloning to an animal cell expression vector

The DNA fragment amplified in 1) was fractionated by 1% agarose gel electrophoresis. After staining the gel with ethidium bromide, the gel containing the desired band observed under ultraviolet irradiation was cut out. The extraction of the DNA fragment from the agarose gel and purification were carried out using a GENECLAN II Kit (Bio101 Co.)

The extracted and purified DNA fragment was subcloned to animal cell expression vector pTARGET (Promega Co.) The ligation solution used was a Takara Ligation Kit Ver. 2 (Takara Shuzo), and the reaction was carried out with the following composition at 16°C for 1.5 h.

Extracted and purified DNA fragment	1 μ L (50 ng)
PTARGET	1 μ L (10 ng)
Water	3 μ L
<u>Ligation solution</u>	<u>5 μL</u>
Total	10 μ L

The reaction solution after the above reaction was used to transform the *E. coli* K12 strain DH5. The transformant was inoculated on an LB agar medium containing 50 μ g/mL of ampicillin (Amp), 40 μ g/mL of 5-bromo-4-chloro-3-indolyl- β -D-galactoside (IPTG) [sic; isopropyl- β -D-thiogalactopyranoside] and 100 μ M of isopropyl β -D-thiogalactopyranoside

(X-gal) [sic; 5-bromo-4-chloro-3-indolyl- β -D-galactopyranoside] * and cultivated overnight at 37°C.

Each colony that developed on the above plate was inoculated in 10 mL of an LB liquid medium containing 50 μ g/mL of Amp, cultivation was carried out overnight at 37°C, the biomass was collected by centrifugation, and subsequently the recombinant DNA was purified using a QIAprep Spin Plasmid Miniprep Kit (Qiagen Co.) to obtain pTARGETpgth.

3) Determination of the base sequence of the inserted cDNA

The base sequence determination was carried out using a DNA sequencer (ABI Co., Model PRISM377) and the dye terminator method, and the whole base sequence of the two chains was determined using the primer walking method. The clone was found to contain all of the region between sequences 1 and 2 among sequence No. 3 confirming that the desired gene pTARGETpgth had been cloned.

Application Example 4

Insertion into CHOK1 cells and stable transformant preparation

The recombinant DNA, pTARGETpgth, prepared in Application Example 2 has a CMV promoter upstream of pgth, and if it is inserted into an animal cell, the expression of pgth is possible.

CHOK1 cells were cultured in 60 mm diameter plastic Petri dishes. As the culture medium, Ham F-12 (Gibco, called growth medium, below) containing 10% fetal bovine serum (Dainippon Seiyaku), 50 U/mL of penicillin and 50 μ g/mL of streptomycin was used, and culture was carried out at 37°C in the presence of 5% CO₂. When the cell density was 50%, LIPOFECTAMINE reagent (Gibco) containing pTARGETpgth prepared in Application Example 2 was added in a layer over the cells, incubated for 6 h, and, after replacement with the growth medium, culture was continued for 48 h. After dispersing the cells with trypsin, the cell suspension was placed in a 60 mm diameter plastic Petri dish, and culture was carried out for 24 h. After removing the culture medium, it was replaced by growth medium containing G418 (Gibco, final concentration of 500 μ g/mL). The G418 medium was changed every 3 days and culture continued for 2 weeks. When the cell colonies were observable with the naked eye, 3 colonies were isolated using stainless steel cups. As a control, only the pTARGET vector (Promega Co.) was inserted into CHOK1 cells by carrying out the same procedures as those described above to isolate a stable transformant.

* [Editor's note: The compound names and abbreviations are so garbled in the original text that it is impossible to be certain whether it should be 40 μ g/mL IPTG and 100 μ M X-gal, or vice-versa.]

washed with a suitable buffer solution containing bovine serum albumin, and culture was continued for 20 min using a buffer solution containing (^3H)-labeled PGE2 (Amersham Co.). After washing the cells, they were recovered, and the radioactivity taken up was measured. As a result, the prostaglandin transport activity of the CHO κ 1 cells with pgth inserted was statistically significantly higher than that of the CHO κ 1 cells with only the control vector inserted.

Application Example 6

Expression of pgth mRNA in human macrophages loaded with oxidized LDL

1) Preparation of human macrophages loaded with oxidized LDL and normal monocyte cDNA

Normal monocyte cDNA was prepared using RNA prepared with Trizol (Gibco BRL Co.) from CD14-positive monocytes from human peripheral blood as a template and the SuperScript II reverse transcriptase kit (Gibco BRL). Human macrophages loaded with oxidized LDL were prepared by culturing normal monocytes in a RPMI-1640 medium (Dainippon Seiyaku) containing 20% AB serum and antibiotics for 14 days, adding human LDL oxidized with copper sulfate using conventional procedures (oxidized LDL) in the final concentration of 40 μmL [sic; dimension incorrect] and continuing culture for 24 h. A method similar to that used for normal monocytes was used to prepare cDNA.

2) Confirmation of pgth mRNA expression by the RT-PCR method

Oligonucleotides (following sequence 3) having a sequence contained in sequence No. 2 and oligonucleotides (following sequence 4) having the sequence of the reverse complementary strand were respectively synthesized using a DNA synthesizer (ABI Co., Model 380B).

Sequence 3

5-GCTCCTGCCCATTGGACGGCTTTAACC-3

Sequence 4

5-TCACACTCGGGAATCCTCTGGCTTC-3

The cDNA prepared in (1) was used as a template, the oligonucleotides with sequences 3 and 4 were used as primers, and the following PCR procedures were carried out using a Takara LA PCR kit Ver. 2 and the PCR thermal cycler MP (Takara Shuzo).

cDNA	2 μL (40 ng)
10X PCR buffer (containing 25 mM Mg^{++})	1.5 μL
2.5 mM dNTP	2.4 μL

10 μ M Sequence 3	0.4 μ L
10 μ M Sequence 4	0.4 μ L
Water	10.15 μ L
LA Taq polymerase	0.15 μ L
Total amount	15 μ L

The PCR cycle was carried out by holding at 94°C for 5 min, carrying out the reaction at 94°C for 1 min, holding at 58°C for 1 min, furthermore at 72°C for 1 min, and repeating 30 times. The PCR reaction mixture was fractionated using 1% agarose gel electrophoresis. After staining the gel with ethidium bromide, the ultraviolet irradiation was carried out to detect an amplified band at about 500 bp. Similarly, the glyceraldehyde 3-phosphate dehydrogenase gene amplified primer (G3PDH, Clontech Co.) was used as the standard cDNA for PCR testing. As a result, the expression of pgth mRNA was strongly induced in the macrophages loaded with oxidized LDL, as shown in Figure 3.

Normal monocytes, macrophages loaded with oxidized LDL or equivalent cultured cells may be cultured with a test compound added, and subsequently the change in the PGTH mRNA may be measured by the method described above to screen any substance controlling PGTH mRNA expression.

Brief description of the figures

Figure 1 shows comparison of amino acid sequence homology between hPGT and the PGTH of this invention.

Figure 2 shows the results of SDS-PAGE of PGTH expressed using the in vitro translation method using pgth.

Figure 3 shows the results of detection of mRNA for the expression of pgth in human macrophages loaded with oxidized LDL using the RT-PCR method. In the figure o shows the results for human macrophages loaded with oxidized LDL, and m shows the results for normal human monocytes.

Claims

(1) A protein of the following (a) or (b).

(a) Protein comprising the amino acid sequence of sequence No. 1

(b) Protein comprising an amino acid sequence with one or more amino acids deleted, substituted or added to the amino acid sequence of sequence No. 1, and, at the same time, having a prostaglandin transport activity.

(2) DNA of the following (a) or (b).

(a) DNA comprising the base sequence of sequence No. 2

(b) DNA which can be hybridized with the DNA of sequence No. 2 under stringent conditions and at the same time, encodes a protein having a prostaglandin transport activity.




Figure 1

Replacement Sheet (Regulation 26)




Figure 2




Figure 3

SEQUENCE LISTING

<110> TAISHO PHARMACEUTICAL CO., Ltd.

<120> Prostaglandin

<130> P487

<150> JP10-227723

<151> 1998-08-12

<160> 3

<210> 1

<211> 709

<212> PRT

<213> Homo sapience

<400> 1

```

Met Gly Pro Arg Ile Gly Pro Ala Gly Glu Val Pro Gln Val Pro
      5                      10                      15
Asp Lys Glu Thr Lys Ala Thr Met Gly Thr Glu Asn Thr Pro Gly
      20                      25                      30
Gly Lys Ala Ser Pro Asp Pro Gln Asp Val Arg Pro Ser Val Phe
      35                      40                      45
His Asn Ile Lys Leu Phe Val Leu Cys His Ser Leu Leu Gln Leu
      50                      55                      60
Ala Gln Leu Met Ile Ser Gly Tyr Leu Lys Ser Ser Ile Ser Thr
      65                      70                      75
Val Glu Lys Arg Phe Gly Leu Ser Ser Gln Thr Ser Gly Leu Leu
      80                      85                      90
Ala Ser Phe Asn Glu Val Gly Asn Thr Ala Leu Ile Val Phe Val
      95                      100                     105
Ser Tyr Phe Gly Ser Arg Val His Arg Pro Arg Met Ile Gly Tyr
      110                     115                     120
Gly Ala Ile Leu Val Ala Leu Ala Gly Leu Leu Met Thr Leu Pro
      125                     130                     135
His Phe Ile Ser Glu Pro Tyr Arg Tyr Asp Asn Thr Ser Pro Glu
      140                     145                     150
Asp Met Pro Gln Asp Phe Lys Ala Ser Leu Cys Leu Pro Thr Thr
      155                     160                     165
Ser Ala Pro Ala Ser Ala Pro Ser Asn Gly Asn Cys Ser Ser Tyr
      170                     175                     180
Thr Glu Thr Gln His Leu Ser Val Val Gly Ile Met Phe Val Ala
      185                     190                     195

```

Gln Thr Leu Leu Gly Val Gly Gly Val	Pro Ile Gln Pro Phe Gly	
200	205	210
Ile Ser Tyr Ile Asp Asp Phe Ala His	Asn Ser Asn Ser Pro Leu	
215	220	225
Tyr Leu Gly Ile Leu Phe Ala Val Thr	Met Met Gly Pro Gly Leu	
230	235	240
Ala Phe Gly Leu Gly Ser Leu Met Leu	Arg Leu Tyr Val Asp Ile	
245	250	255
Asn Gln Met Pro Glu Gly Gly Ile Ser	Leu Thr Ile Lys Asp Pro	
260	265	270
Arg Trp Val Gly Ala Trp Trp Leu Gly	Phe Leu Ile Ala Ala Gly	
275	280	285
Ala Val Ala Leu Ala Ala Ile Pro Tyr	Phe Phe Phe Pro Lys Glu	
290	295	300
Met Pro Lys Glu Lys Arg Glu Leu Gln	Phe Arg Arg Lys Val Leu	
305	310	315
Ala Val Thr Asp Ser Pro Ala Arg Lys	Gly Lys Asp Ser Pro Ser	
320	325	330
Lys Gln Ser Pro Gly Glu Ser Thr Lys	Lys Gln Asp Gly Leu Val	
335	340	345
Gln Ile Ala Pro Asn Leu Thr Val Ile	Gln Phe Ile Lys Val Phe	
350	355	360
Pro Arg Val Leu Leu Gln Thr Leu Arg	His Pro Ile Phe Leu Leu	
365	370	375
Val Val Leu Ser Gln Val Cys Leu Ser	Ser Met Ala Ala Gly Met	
380	385	390
Ala Thr Phe Leu Pro Lys Phe Leu Glu	Arg Gln Phe Ser Ile Thr	
395	400	405
Ala Ser Tyr Ala Asn Leu Leu Ile Gly	Cys Leu Ser Phe Pro Ser	
410	415	420
Val Ile Val Gly Ile Val Val Gly Gly	Val Leu Val Lys Arg Leu	
425	430	435
His Leu Gly Pro Val Gly Cys Gly Ala	Leu Cys Leu Leu Gly Met	
440	445	450
Leu Leu Cys Leu Phe Phe Ser Leu Pro	Leu Phe Phe Ile Gly Cys	
455	460	465
Ser Ser His Gln Ile Ala Gly Ile Thr	His Gln Thr Ser Ala His	
470	475	480
Pro Gly Leu Glu Leu Ser Pro Ser Cys	Met Glu Ala Cys Ser Cys	
485	490	495
Pro Leu Asp Gly Phe Asn Pro Val Cys	Asp Pro Ser Thr Arg Val	
500	505	510
Glu Tyr Ile Thr Pro Cys His Ala Gly	Cys Ser Ser Trp Val Val	
515	520	525
Gln Asp Ala Leu Asp Asn Ser Gln Val	Phe Tyr Thr Asn Cys Ser	
530	535	540
Cys Val Val Glu Gly Asn Pro Val Leu	Ala Gly Ser Cys Asp Ser	
545	550	555
Thr Cys Ser His Leu Val Val Pro Phe	Leu Leu Leu Val Ser Leu	
560	565	570

Gly Ser Ala Leu	Ala Cys Leu Thr His Thr Pro Ser Phe Met Leu	
575	580	585
Ile Leu Arg Gly	Val Lys Lys Glu Asp Lys Thr Leu Ala Val Gly	
590	595	600
Ile Gln Phe Met	Phe Leu Arg Ile Leu Ala Trp Met Pro Ser Pro	
605	610	615
Val Ile His Gly	Ser Ala Ile Asp Thr Thr Cys Val His Trp Ala	
620	625	630
Leu Ser Cys Gly	Arg Arg Ala Val Cys Arg Tyr Tyr Asn Asn Asp	
635	640	645
Leu Leu Arg Asn	Arg Phe Ile Gly Leu Gln Phe Phe Phe Lys Thr	
650	655	660
Gly Ser Val Ile	Cys Phe Ala Leu Val Leu Ala Val Leu Arg Gln	
665	670	675
Gln Asp Lys Glu	Ala Arg Thr Lys Glu Ser Arg Ser Ser Pro Ala	
680	685	690
Val Glu Gln Gln	Leu Leu Val Ser Gly Pro Gly Lys Lys Pro Glu	
695	700	705
Asp Ser Arg Val		
709		

<210> 2
 <211> 2130
 <212> DNA
 <213> Homo sapiens

<400> 2

10	20	30	40	50	60	
algggaccca	ggala	gggccc	agcggg	lgag	giacccagg	laccagacaa ggaaaccaa 60
gccacaalgg	gcacagaaa	cacaccigga	ggcaaagcca	gcccagaccc	tcaggacgig	120
cggccaagig	tgltccataa	catraagcig	ttcgltcigt	gccacagcct	gcigcagcig	180
ggcgagctca	tgatctccgg	ctacctaag	agctccatct	ccacagigga	gaagcgcttc	240
ggcctctcca	gccagacgic	ggggcigcig	gctctctcca	acgaggiggg	gaacacagcc	300
ttgattgigt	ttgtgagcta	tttgggcagc	cgggtgcacc	gaccccgaa	gattggctat	360
ggggctatcc	ttgtggccct	ggcgggccc	ctcatgac	ctccgcactt	catctcggag	420
ccataccgct	acgacaacac	cagcccigag	gatalgccac	aggacttcaa	ggcttccctg	480
tgccigccca	caacctcggc	cccagccctg	gccccctcca	atggcaactg	ctcaagctac	540
acagaaaccc	agcatctgag	tggtggggg	atcatgttgc	tggcacagac	ccigctgggc	600

giggcgggg igcccaica gccctllggc atctctaca tcatgactt igcccacaac 660
agcaaclegc cccctiacct cgggaltcig lllgcagiga ccatgaiggg gccaggccig 720
gcclllgggc lgggcagccf catgcigcgc cltlatlggg acallaacca galgccagaa 780
ggtgglatca gccigaccat aaaggacccc cgaigggigg glgcciglg gclggglttc 840
clcatcgcig ccggigcagi ggccllggt gccatcccc acttctctt ccccaaggaa 900
algcceaagg aaaaacgiga gcltcaglll cggcgaaagg tcltagcagi cacagactca 960
cclgccagga agggcaagga ctctccctt aagcagagcc clggggagtc cacaagaag 1020
caggatggcc tagtcagat tgcaccaaac ctgactgiga tccagltcat taaaglttc 1080
cccagggtgc lgcigcagac cctacgccac cccatcttcc tgcigggigt cclgtcccag 1140
glatgcltgi calccaiggc tgggggaig gccaccttcc lgcceaagtl cclggagcgc 1200
cagltttcca tcaagccct clacgccaac ctgtcaltg gclgccttc ctcccttcg 1260
glcatcgtgg gcatcgtgtt gggiggcgt ctggtcaagc ggctccacct gggccclgtg 1320
ggatgcgglg ccclllgct gclggggaig ctgclgtgcc tcltcltcag cclgccgtc 1380
tclttlatcg gclgtccag ccaccagat gggggcaica cacaccagac caglgccac 1440
cclgggcgtg agclgtctcc aagclgcalg gaggcctgt cclgccat ggacggcttl 1500
aacclgtct ggcaccccag cactcgtgtg gaalacaica caccclgcca cgcaggctgc 1560
tcaagclggg lggltccagga tgccltgac aacagccagg tllcttacc caacigcagc 1620
tgcgtgglg agggcaacc cglgtggca ggalcctgc actcaacgtg cagccatctg 1680
glgglgccct tclgtctctt ggtcagccgt ggcctggccc tggccgtct caccacaca 1740
ccccttca tgcaltctt aagaggagtg aagaaagaag acaagactt ggcgtgggc 1800
altcagltca tglcttgag gatlllgcc tggatgccc gccccgtgat ccacggcagc 1860
gccatcgaca ccaccigtgt gcactggcc ctgagctgt ggcgtcgagc tgcgtctgc 1920
tactacaata atgacctgt ccgaaaccgg ttcactggcc tccagltct cltcaaaaca 1980
ggtlctgiga tclgtctgc cltaglttg gclgtctga ggcagcagga caaagaggca 2040
aggaccaaaag agagcagat cagccctgcc glagagcagc aatlgctagt gtcggggcca 2100

gggaagaagc cagaggattc ccgagtgga

2130

<210> 3
 <211> 4083
 <212> DNA
 <213> Homo sapiens

<400> 3

aagtagacca gggagacaaa cacttggaga tacttggggc tgagttag caagactccc 60
 taaccttggt cttgacaagt ctgattgctt gttgggccc agaagaactg acccctgttc 120
 tggagctccc accgttatgt catccctgct gttggctacc tgcctgtgtc tccaggagcc 180
 cctgagaaga ttgtccctct cttccctgct aagctccagg tcttgagatt gaattagggg 240
 ctggagctca ctgcactcca gcagtc 266

atg gga ccc agg ata ggg cca gcg ggt gag gta ccc cag gla cca 311
 Met Gly Pro Arg Ile Gly Pro Ala Gly Glu Val Pro Gln Val Pro
 5 10 15
 gac aag gaa acc aaa gcc aca atg ggc aca gaa aac aca cct gga 356
 Asp Lys Glu Thr Lys Ala Thr Met Gly Thr Glu Asn Thr Pro Gly
 20 25 30
 ggc aaa gcc agc cca gac cct cag gac gtt cgg cca agt gtt ttc 401
 Gly Lys Ala Ser Pro Asp Pro Gln Asp Val Arg Pro Ser Val Phe
 35 40 45
 cat aac atc aag ctg ttc gtt ctg tgc cac agc ctg ctg cag ctg 446
 His Asn Ile Lys Phe Val Leu Cys His Ser Leu Leu Gln Leu
 50 55 60
 gcg cag ctg atg atc tcc gcc tac cta aag agc tcc atc tcc aca 491
 Ala Gln Leu Met Ile Ser Gly Tyr Leu Lys Ser Ser Ile Ser Thr
 65 70 75
 gtt gag aag cgc ttc gcc ctg tcc agc cag acg tgg ggg ctg ctg 536
 Val Glu Lys Arg Phe Gly Leu Ser Ser Gln Thr Ser Gly Leu Leu
 80 85 90
 gcc tcc ttc aac gag gtt ggg aac aca gcc ttg att gtt ttt gtt 581
 Ala Ser Phe Asn Glu Val Gly Asn Thr Ala Leu Ile Val Phe Val
 95 100 105
 agc tat ttt ggc agc cgg gtt cac cga ccc cga atg att ggc tat 626
 Ser Tyr Phe Gly Ser Arg Val His Arg Pro Arg Met Ile Gly Tyr
 110 115 120
 ggg gct atc ctt gtt gcc ctg gcg ggc ctg ctg atg act ctg ccg 671
 Gly Ala Ile Leu Val Ala Leu Ala Gly Leu Leu Met Thr Leu Pro
 125 130 135
 cac ttc atc tgg gag cca tac cgc tac gac aac acc agc cct gag 716
 His Phe Ile Ser Glu Pro Tyr Arg Tyr Asp Asn Thr Ser Pro Glu
 140 145 150

gat atg cca cag gac ttc aag gct tcc ctg tgc ctg ccc aca acc	761
Asp Met Pro Gln Asp Phe Lys Ala Ser Leu Cys Leu Pro Thr Thr	
155 160 165	
tcg gcc cca gcc tcg gcc ccc tcc aat ggc aac tgc tca agc tac	806
Ser Ala Pro Ala Ser Ala Pro Ser Asn Gly Asn Cys Ser Ser Tyr	
170 175 180	
aca gaa acc cag cal ctg agt gtg gtg ggg atc atg ttc gtg gca	851
Thr Glu Thr Gln His Leu Ser Val Val Gly Ile Met Phe Val Ala	
185 190 195	
cag acc ctg ctg ggc gtg ggc ggg gtg ccc att cag ccc ttt ggc	896
Gln Thr Leu Leu Gly Val Gly Gly Val Pro Ile Gln Pro Phe Gly	
200 205 210	
atc tcc tac atc gat gac ttt gcc cac aac agc aac tcg ccc ctg	941
Ile Ser Tyr Ile Asp Asp Phe Ala His Asn Ser Asn Ser Pro Leu	
215 220 225	
tac ctg ggg atc ctg ttt gca gtg acc atg atg ggg cca ggc ctg	986
Tyr Leu Gly Ile Leu Phe Ala Val Thr Met Met Gly Pro Gly Leu	
230 235 240	
gcc ttt ggg ctg ggc agc ctg atg ctg cgc ctt tat gtg gac att	1031
Ala Phe Gly Leu Gly Ser Leu Met Leu Arg Leu Tyr Val Asp Ile	
245 250 255	
aac cag atg cca gaa ggt ggt atc agc ctg acc ata aag gac ccc	1076
Asn Gln Met Pro Glu Gly Gly Ile Ser Leu Thr Ile Lys Asp Pro	
260 265 270	
cga tgg gtg ggt gcc tgg tgg ctg ggt ttc ctg atc gct gcc ggt	1121
Arg Trp Val Gly Ala Trp Trp Leu Gly Phe Leu Ile Ala Ala Gly	
275 280 285	
gca gtg gcc ctg gct gcc atc ccc tac ttc ttc ttc ccc aag gaa	1166
Ala Val Ala Leu Ala Ala Ile Pro Tyr Phe Phe Phe Pro Lys Glu	
290 295 300	
atg ccc aag gaa aaa cgt gag ctt cag ttt cgg cga aag gtc tta	1211
Met Pro Lys Glu Lys Arg Glu Leu Gln Phe Arg Arg Lys Val Leu	
305 310 315	
gca gtc aca gac tca cct gcc agg aag ggc aag gac tct ccc tct	1256
Ala Val Thr Asp Ser Pro Ala Arg Lys Gly Lys Asp Ser Pro Ser	
320 325 330	
aag cag agc cct ggg gag tcc acg aag aag cag gat ggc cta gtc	1301
Lys Gln Ser Pro Gly Glu Ser Thr Lys Lys Gln Asp Gly Leu Val	
335 340 345	
cag att gca cca aac ctg act gtg atc cag ttc att aaa gtc ttc	1346
Gln Ile Ala Pro Asn Leu Thr Val Ile Gln Phe Ile Lys Val Phe	
350 355 360	
ccc agg gtg ctg ctg cag acc cta cgc cac ccc atc ttc ctg ctg	1391
Pro Arg Val Leu Leu Gln Thr Leu Arg His Pro Ile Phe Leu Leu	
365 370 375	
gtg gtc ctg tcc cag gta tgc ttg tca tcc atg gct gcg ggc atg	1436
Val Val Leu Ser Gln Val Cys Leu Ser Ser Met Ala Ala Gly Met	
380 385 390	
gcc acc ttc ctg ccc aag ttc ctg gag cgc cag ttt tcc atc aca	1481
Ala Thr Phe Leu Pro Lys Phe Leu Glu Arg Gln Phe Ser Ile Thr	

	395	400	405	
gcc tcc tac gcc aac ctc ctc atc gcc tgc ctc tcc ttc cct tgc				1526
Ala Ser Tyr Ala Asn Leu Leu Ile Gly Cys Leu Ser Phe Pro Ser				
	410	415	420	
gtc atc glg gcc atc glg glg ggt gcc gtc ctc gtc aag cgg ctc				1571
Val Ile Val Gly Ile Val Val Gly Gly Val Leu Val Lys Arg Leu				
	425	430	435	
cac ctc gcc cct gtg gga tgc ggt gcc ctt tgc ctc ctc ggg atg				1616
His Leu Gly Pro Val Gly Cys Gly Ala Leu Cys Leu Leu Gly Met				
	440	445	450	
ctg ctc tgc ctc ttc ttc agc ctc ccg ctc ttc ttt atc gcc tgc				1661
Leu Leu Cys Leu Phe Phe Ser Leu Pro Leu Phe Phe Ile Gly Cys				
	455	460	465	
tcc agc cac cag att gcg gcc atc aca cac cag acc agt gcc cac				1706
Ser Ser His Gln Ile Ala Gly Ile Thr His Gln Thr Ser Ala His				
	470	475	480	
cct ggg ctc gag ctc tct cca agc tgc atg gag gcc tgc tcc tgc				1751
Pro Gly Leu Glu Leu Ser Pro Ser Cys Met Glu Ala Cys Ser Cys				
	485	490	495	
cca ttg gac gcc ttt aac cct gtc tgc gac ccc agc act cgt glg				1796
Pro Leu Asp Gly Phe Asn Pro Val Cys Asp Pro Ser Thr Arg Val				
	500	505	510	
gaa tac atc aca ccc tgc cac gca gcc tgc tca agc tgg glg gtc				1841
Glu Tyr Ile Thr Pro Cys His Ala Gly Cys Ser Ser Trp Val Val				
	515	520	525	
cag gat gct ctc gac aac agc cag gtt ttc tac acc aac tgc agc				1886
Gln Asp Ala Leu Asp Asn Ser Gln Val Phe Tyr Thr Asn Cys Ser				
	530	535	540	
tgc glg glg gag gcc aac ccc glg ctc gca gga tcc tgc gac tca				1931
Cys Val Val Glu Gly Asn Pro Val Leu Ala Gly Ser Cys Asp Ser				
	545	550	555	
acg tgc agc cat ctc glg glg ccc ttc ctc ctc ctc gtc agc ctc				1976
Thr Cys Ser His Leu Val Val Pro Phe Leu Leu Leu Val Ser Leu				
	560	565	570	
ggc tgc gcc ctc gcc tgt ctc acc cac aca ccc tcc ttc atg ctc				2021
Gly Ser Ala Leu Ala Cys Leu Thr His Thr Pro Ser Phe Met Leu				
	575	580	585	
atc cta aga gga glg aag aaa gaa gac aag act ttg gct glg gcc				2066
Ile Leu Arg Gly Val Lys Lys Glu Asp Lys Thr Leu Ala Val Gly				
	590	595	600	
atc cag ttc atg ttc ctc agg att ttg gcc tgg atg ccc agc ccc				2111
Ile Gln Phe Met Phe Leu Arg Ile Leu Ala Trp Met Pro Ser Pro				
	605	610	615	
gtg atc cac gcc agc gcc atc gac acc acc tgt glg cac tgg gcc				2156
Val Ile His Gly Ser Ala Ile Asp Thr Thr Cys Val His Trp Ala				
	620	625	630	
ctg agc tgt ggg cgt cga gct glc tgt cgc tac tac aat aat gac				2201
Leu Ser Cys Gly Arg Arg Ala Val Cys Arg Tyr Tyr Asn Asn Asp				
	635	640	645	
ctg ctc cga aac cgg ttc atc gcc ctc cag ttc ttc ttc aaa aca				2246

Leu	Leu	Arg	Asn	Arg	Phe	Ile	Gly	Leu	Gln	Phe	Phe	Phe	Lys	Thr		
			650						655					660		
ggg	tcg	gig	atc	lgc	ttc	gcc	tta	gtt	tig	gct	gtc	cig	agg	cag	2291	
Gly	Ser	Val	Ile	Cys	Phe	Ala	Leu	Val	Leu	Ala	Val	Leu	Arg	Gln		
			665						670					675		
cag	gac	aaa	gag	gca	agg	acc	aaa	gag	agc	aga	icc	agc	ccg	gcc	2336	
Gln	Asp	Lys	Glu	Ala	Arg	Thr	Lys	Glu	Ser	Arg	Ser	Ser	Pro	Ala		
			680						685					690		
gta	gag	cag	caa	tig	cta	gig	tcg	ggg	cca	ggg	aag	aag	cca	gag	2381	
Val	Glu	Gln	Gln	Leu	Leu	Val	Ser	Gly	Pro	Gly	Lys	Lys	Pro	Glu		
			695						700					705		
gal	icc	cga	gtg	lga											2396	
Asp	Ser	Arg	Val													
			709													

gcgtcttgg ggccccacct ggccaagagt agcagccaca gcagtaacct ccttgagacc 2456

ttggcccaag attgggggtc aagagcccig tgttccattc tggctccccc actaaatggc 2516

tgtgtgacct caggcaagac attgataccc tctcagccct tgccttgctag tctgaaccaa 2576

agagttgttt gggcatttgc tgtgttggcc attcttgag caagagggtc ttcttccccc 2636

ttccccagc cagccagctg tcttggggcc aggccttccc gggtagaaag aagtaacct 2696

ttccctgggg ccttaggata gcaaagtgag ccatagtggg ccaggctgcc ctccatgctg 2756

ggccccagcc caggctcgca ctggccggga tcacctctt ttagccctag ccattctctg 2816

tcaggtagga atgaacttgc cagccctcag gctcgttcag ctatgacct ctgtgcgggc 2876

agggtacact cagctctccc ccccaacctc agcagccctt aagaagtgc ccttgggcgc 2936

ccccggagg cagagcacg agcaggacc tgggtagact cccacaggga ggacggagct 2996

ggcctcagga gtgggacacc cagacttggc agggccctca agaggccctg gtggggggcc 3056

caggaatccc tagctgaagc ggggagacac actctccatc tcaggaaatt ctagcccttg 3116

ccctcagga gccacggctg agggtaggc ccaacacctg ccttagggcc ctgggtgggc 3176

aagcttgggc cctggggtag ggaggagac tcaggccac acttgggtat ttctaat 3236

cagacaaaca cacactcagc gcgcactcac tgattctac acatggccaa gattcacac 3296

atgtgaccag ggccaccaaa agtcccttgc acccttggga ctaggatccc aattctctta 3356

tttctctctg ggtgcttggg tctgtgtcac ctggggcagt gtggataatg tttagttctg 3416

tgacactgtt ttttgggggt ggcacctgtt tctccgatgc ctgggcctgt gtcaggccca 3476

ggactgtagt gcigggagca glaaagctca gcicigigia algaglgalg clatggcctg 3536
ctcgtgicct atgalccaat cctilicctac alcageccct gllltglll alggctagtc 3596
ttatciggcc tggllatitc ctgcgggga ggagagggt tgciaalcig ctcccagccc 3656
aacctallac caccaccct cgcigggacc tactgcicgg gaggcagcag acaggagacc 3716
accagcagtg gcttcciggc cctigcicgg gggiggggg aagciggggg cacaigtggc 3776
cctigccitc tgagcagctc ccaglgccag ggctllgaga ctileccaca tgalaaaaga 3836
aaaggagggt acagaagtic caatccctt ttatlltgc tggllggial ctglaaatgt 3896
ttaataaata tcigagcaig tatctatcaa cgccaagaal ttcaaagctt ccttcaacaa 3956
tatgaggctt ttaggatgtt tatattcctt catccctctt glltcccagg ttllgcaggg 4016
aaaaaaagtc tggaaatlala galacagctt attattaaat llttctctgc ataaaaaaaa 4076
aaaaaaa 4083

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04352

A. CLASSIFICATION OF SUBJECT MATTER

Int.Cl.⁶ C07K 14/47, C12N 15/12//C12N 5/10, C12P 21/02,
(C12P 21/02, C12R 1:91)

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

Int.Cl.⁶ C07K 14/47, C12N 15/12, C12N 5/10, C12P 21/02,

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)
SwissProt/PIR/GeneSeq, Genbank/EMBL/DDBJ/GeneSeq,
WPI (DIALOG), BIOSIS (DIALOG)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	US, 5792851, A (Albert Einstein College of Medicine of Yeshiva University, a Division of Yeshiva University) 11 August, 1998 (11.08.98) (Family: none)	1, 2
Y	Journal of Clinical Investigation, Vol. 98, no. 5 (1996) Lu Run, et al., "Cloning, in vitro expression, and tissue distribution of a human prostaglandin transporter cDNA (hPGT)" see p.1142-1149, (1996)	1, 2
Y	Biochemical and Biophysical Research Communications, Vol. 246, No. 3, (May 29, 1998), Lu Run, et al., "Molecular cloning of the gene for human prostaglandin transporter hPGT: Gene organization, promoter activity and chromosomal localization", see p.805-812,	1, 2

☒ Further documents are listed in the continuation of Box C.

☐ See patent family annex.

* Special categories of cited documents:
"A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Date of the actual completion of the international search
09 November, 1999 (09.11.99)

Date of mailing of the international search report
24 November, 1999 (24.11.99)

Name and mailing address of the ISA/
Japanese Patent Office

Authorized officer

Facsimile No.

Telephone No.

INTERNATIONAL SEARCH REPORT

International application No.

PCT/JP99/04352

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	Science, vol.268, No.5212, (1995), Kanai Naoaki et al., "Identification and Characterization of a prostaglandin transporter ", see p.866-869,	1-2